



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SOLUTION OF PROBLEMS IN NUMBER FOUR.


SOLUTIONS of problems in No. 4 have been received as follows:

From R. J. Adcock, 120 and 123; Marcus Baker, 119, 122 and 123; Dr. H. Eggers, 122; Edgar Frisby, 121; Henry Heaton, 119, 120, 121*, 122, 123 and 124; Prof. H. T. J. Ludwick, 121*; L. Regan, 119 and 122; E. B. Seitz, 119, 122, 123 and 124.

119. "If from one of the extremities, B , of the diameter, AB , of a given circle, any chord, BD , be drawn, and from the other extremity, A , and the centre C , two lines, AE , CE , be drawn to any point, E , in the circumference, cutting said, chord in the points F and G ; then, $GF : GD :: (BF)^2 : (BA)^2$. Required the demonstrator."

DEMONSTRATION BY E. B. SEITZ, GREENVILLE, OHIO.

Draw GH parallel to BA ; join DE and DH . The similar triangles DEF and HGF give $DF : HF :: EF : GF$, or $DF : EF :: HF : GF$; hence the triangles DFH and EFG are similar; therefore $\angle HDG = \angle GEH = \angle A$, and consequently the triangles ABF and DGH are similar.



The similar triangles ABF , HGF and DGH give $GF : GH :: BF : BA$, and $GH : GD :: BF : BA$. Multiplying these proportions together, we have $GF : GD :: BF^2 : BA^2$.

120. [As the solutions of this question, that have been submitted, are only forms from which the numerical value of x and y can be found by approximation, and as the answer which was sent with the question by Dr. Oliver, the proposer, cannot be verified by assigning particular values to x and y , and is therefore erroneous, it is not likely that a general solution of the question can be obtained.]

121 "In a spherical triangle are given the sum of each angle and the side opposite, to solve the triangle."

SOLUTION BY EDGAR FRISBY, ESQ., NAVAL OBSERVATORY, WASH., D.C.

We have

$$(1) \frac{\sin A - \sin a}{\sin A + \sin a} = \frac{\sin B - \sin b}{\sin B + \sin b} = \frac{\sin C - \sin c}{\sin C + \sin c} = \frac{\tan \frac{1}{2}(A-a)}{\tan \frac{1}{2}(A+a)} \&c. = x \text{ say.}$$

(2) $\tan \frac{1}{2}(A-a) = lx, \tan \frac{1}{2}(B-b) = mx, \tan \frac{1}{2}(C-c) = nx$ if $\tan \frac{1}{2}(A+a) = l$ &c.

$$(3) \begin{cases} \sin A = [\tan \frac{1}{2}(A+a) + \tan \frac{1}{2}(A-a)] \cos \frac{1}{2}(A+a) \cos \frac{1}{2}(A-a) \\ \quad = \frac{l(1+x)}{\sqrt{[(1+l^2)(1+l^2x^2)]}}; \cos A = \frac{1+l^2x}{\sqrt{[(1+l^2)(1+l^2x^2)]}} \\ \sin a = \frac{l(1-x)}{\sqrt{[(1+l^2)(1+l^2x^2)]}}; \cos a = \frac{1+l^2x}{\sqrt{[(1+l^2)(1+l^2x^2)]}} \end{cases}$$

and similar expressions for B, b, C and c ; substituting these values in Cagnoli's equation

$$(4) \quad \sin B \sin C - \sin b \sin c = \cos B \cos C \cos a + \cos b \cos c \cos A$$

we have

$$(5) \quad \frac{mn[(1+x)^2 - (1-x)^2]}{\sqrt{[(1+m^2)(1+n^2)(1+m^2x^2)(1+n^2x^2)]}} \\ = \frac{(1-m^2x)(1-n^2x)(1+l^2x) - (1+m^2x)(1+n^2x)(1-l^2x)}{\sqrt{[(1+l^2)(1+m^2)(1+n^2)(1+l^2x^2)(1+m^2x^2)(1+n^2x^2)]}}, \text{ or}$$

$$(6) \quad 2mnx\sqrt{[(1+l^2)(1+l^2x^2)]} = 1 + (m^2n^2 - l^2n^2 - l^2m^2)x^2,$$

which, on clearing of radicals, becomes

$$1 - 2(l^2m^2 + m^2n^2 + n^2l^2 + 2l^2m^2n^2)x^2 + (l^4m^4 + m^4n^4 + n^4l^4 - 2l^2m^2n^4 - 2l^2m^4n^2 \\ - 2l^4m^2n^2 - 4l^2m^2n^2)x^4 = 0;$$

dividing by x^4 and solving we have

$$(7) \quad x^{-2} = l^2m^2 + m^2n^2 + n^2l^2 + 2l^2m^2n^2 \pm 2lmn\sqrt{[(1+l^2)(1+m^2)(1+n^2)]}, \text{ or}$$

$$(8) \quad 1 - x^{-2} = (1+l^2)(1+m^2)(1+n^2) - l^2(1+m^2)(1+n^2) - m^2(1+n^2)(1+l^2) \\ - n^2(1+l^2)(1+m^2) \pm 2lmn\sqrt{[(1+l^2)(1+m^2)(1+n^2)]}$$

$$= (1+l^2)(1+m^2)(1+n^2)$$

$$\times \left(1 - \frac{l^2}{1+l^2} - \frac{m^2}{1+m^2} - \frac{n^2}{1+n^2} \mp 2 \frac{l}{\sqrt{(1+l^2)}} \cdot \frac{m}{\sqrt{(1+m^2)}} \cdot \frac{n}{\sqrt{(1+n^2)}} \right) \\ = \frac{1 - \cos^2 \alpha - \cos^2 \beta - \cos^2 \gamma \mp 2 \cos \alpha \cos \beta \cos \gamma}{\sin^2 \alpha \sin^2 \beta \sin^2 \gamma}.$$

If $l = \pm \cot \alpha, m = \pm \cot \beta, n = \pm \cot \gamma$, or $A+a = 180^\circ \mp 2\alpha, B+b = 180^\circ \mp 2\beta, C+c = 180^\circ \mp 2\gamma$, from (2).

In (8) let $x = \cos \varphi$, which is always possible, for from (1) we can show that x always lies between $+1$ and -1 , and equation (8) becomes

$$(9) \quad \tan \varphi = \frac{2\sqrt{[\cos \frac{1}{2}(\alpha+\beta+\gamma) \cdot \cos \frac{1}{2}(\beta+\gamma-\alpha) \cdot \cos \frac{1}{2}(\alpha-\beta+\gamma) \cdot \cos \frac{1}{2}(\alpha+\beta-\gamma)]}}{\sin \alpha \sin \beta \sin \gamma}$$

For the upper sign, by comparing equation (6) with (7) we see that the positive sign only is admissible.

If now we put $\alpha+\beta+\gamma = 2\omega$ this equation becomes

$$\tan \varphi = \frac{+2\sqrt{[\cos \omega \cos (\omega - \alpha) \cos (\omega - \beta) \cos (\omega - \gamma)]}}{\sin \alpha \sin \beta \sin \gamma}$$

which can easily be proved to be always possible. The other root will be

$$\tan \varphi = \frac{+2\sqrt{[-\sin \omega \sin(\omega - \alpha) \sin(\omega - \beta) \sin(\omega - \gamma)]}}{\sin \alpha \sin \beta \sin \gamma},$$

whether this value is real or not can be immediately inferred by inspection; it can be proved that $\alpha, \beta, \gamma, \omega, (\omega - \alpha), (\omega - \beta)$ and $(\omega - \gamma)$ always lie between $\pm 90^\circ$ when the sides and angles are each less than 180° .

Again from (2) $\tan \frac{1}{2}(A - \alpha) = lx = \cot \alpha \cos \varphi$, &c.

We have then $A + \alpha = 180^\circ \pm 2\alpha$, $B + \beta = 180^\circ \pm 2\beta$, $C + \gamma = 180^\circ \pm 2\gamma$, $\alpha + \beta + \gamma = 2\omega$;

$$\tan \varphi = \frac{+2\sqrt{[\cos \omega \cos(\omega - \alpha) \cos(\omega - \beta) \cos(\omega - \gamma)]}}{\sin \alpha \sin \beta \sin \gamma},$$

which is always possible, or

$$\tan \varphi = \frac{+2\sqrt{[-\sin \omega \sin(\omega - \alpha) \sin(\omega - \beta) \sin(\omega - \gamma)]}}{\sin \alpha \sin \beta \sin \gamma},$$

$\tan \frac{1}{2}(A - \alpha) = \cos \varphi \cot \alpha$, $\tan \frac{1}{2}(B - \beta) = \cos \varphi \cot \beta$, $\tan \frac{1}{2}(C - \gamma) = \cos \varphi \cot \gamma$.

It will not make any difference whether we use $A + \alpha = 180^\circ + 2\alpha$ or $A + \alpha = 180^\circ - 2\alpha$ &c., for the sign of $\tan \varphi$ will correspondingly change; $\tan \varphi$ and $\cos \varphi$ must have the same sign.

121*.—“It has been cloudy during the last seven days; what is the probability that it will be cloudy to-morrow?”

SOLUTION BY PROF. H. T. J. LUDWICK, SALISBURY, N. C.

Let the probability that it will be cloudy on any particular day be denoted by x , then $1 - x$ is the probability that it will not be cloudy.

As x may vary from 0 to 1, and since it has been cloudy seven days

∴ $\int_0^1 x^7 dx =$ whole number of possible producing causes

and $\int_0^1 x^8 dx =$ “ “ “ favorable causes.


Therefore the probability that it will be cloudy on the 8th day

$$= \frac{\int_0^1 x^8 dx}{\int_0^1 x^7 dx} = \frac{8}{9}.$$

122. "To inscribe a square in a given quadrilateral."

SOLUTION BY HENRY HEATON, B. S., DES MOINES, IOWA.

Let $ABCD$ be the given quadrilateral. At any convenient distance from the base, as a , draw PQ parallel to the base and cutting AD in P and BC in Q , and lay off PS and QR each equal to a . Draw AS and prolong it to meet BE , drawn perpendicular to AB , in E . Draw BR and prolong it to meet AF , drawn perpendicular to AB , in F , and join EF cutting DC in G . From G draw GO perpendicular to AB cutting BF in M and AE in L , and draw MK and LH parallel to AB . Join GH and GK and draw KI and HI respectively parallel to GH and GK ; then is $GHIK$ the inscribed square required.



Because LO in the triangle EAB equals GM in the triangle EFB , $\therefore GL = MO$. By construction we have $BM : BR :: OM : a :: KM : a$; $\therefore OM = KM$: Also, $AL : AS :: OL : a :: HL : a$; $\therefore OL = HL$. Hence the right-angled triangles GMK and GLH are equal in all their parts and therefore HGK is a right angle and GH IK is a square. Draw IN perpendicular to MK ; then is $IN = GL = MO$. Hence I is on AB and $GHIK$ is the required square.

If the line EF should not intersect CD it is evident there can be no solution; if it should coincide with it, there will be an infinite number.

Problem No. 62 of the ANALYST may be constructed as follows:

Draw BT and AV in the given directions; BT cutting AD in T and AV cutting BC in V , and join TV cutting CD in G . Then is G the vertex of an angle of the required parallelogram.

123. "Solve the equation $\sqrt[2]{x} = a$ and determine what values of a give real roots."

SOLUTION BY MARCUS BAKER, U. S. COAST SURVEY, WASHINGTON, D. C.

Taking the Napierian logarithms of both members we have

$$(\log x) \div x = \log a = b.$$

Put $x = 1 - y$ and substitute for $\log x$ its value from the logarithmic series,

$$\frac{y + \frac{1}{2}y^2 + \frac{1}{3}y^3 + \frac{1}{4}y^4 + \frac{1}{5}y^5 + \frac{1}{6}y^6 + \dots}{1 - y} = -b = c,$$

or, performing the division,

$$x_i = x + \frac{xy^2}{a^2} \dots (1); \quad y_i = -\frac{x^2y}{a^2}; \dots (2)$$

By eliminating x and y from (1) and (2) we get

$$16a^2[x_i + \sqrt{(x_i^2 - 8y_i^2)}] = [3x_i + \sqrt{(x_i^2 - 8y_i^2)}]^3, \dots (3)$$

which is the equation of the curve referred to rectangular coordinates.

From (1) and (2), $dx_i = a^{-2}(3y^2 - a^2)dx$ and $dy_i = a^{-2}(3y^2 - a^2)dy$. Therefore, if z_i = the length of the required curve and z , that of the corresponding arc of the circle, $dz_i = a^{-2}(3y^2 - a^2)dz$ (4)

Put ρ = the radius of curvature of the envelope. Then, as the tangents of the circle and envelope, at their corresponding points, are parallel, it follows that $\rho : a :: dz_i : dz$. $\therefore \rho = a^{-1}(3y^2 - a^2)$. Hence, when $\rho=0$, as at D, E, F, G , $y = a\sqrt{\frac{1}{3}}$, $y_i = \pm \frac{2}{3}a\sqrt{3}$, $x_i = \pm \frac{4}{3}a\sqrt{6}$, and the curve intersects its evolute in the points D, E, F, G , and consists of four branches, of which DAE is the involute of DJ and EJ , DCG , the involute of DI and GI , &c.

From (4) we have $dz_i = a^{-2}(3y^2 - a^2)dz = a^{-1}(3y^2 - a^2)(a^2 - y^2)^{-\frac{1}{2}}dy$; or, because the whole envelope is four times CD plus four times DA ,

$$\begin{aligned} z_i &= \frac{4}{a} \int_{a\sqrt{\frac{1}{3}}}^0 \frac{(3y^2 - a^2)dy}{\sqrt{(a^2 - y^2)}} + \frac{4}{a} \int_0^a \frac{(3y^2 - a^2)dy}{\sqrt{(a^2 - y^2)}} \\ &= 4a(\sqrt{2} - \sin^{-1}\sqrt{\frac{1}{3}}) + a\pi. \end{aligned}$$

If A = the whole area of the envelope, we have

$$\begin{aligned} dA &= y, dx_i = a^{-4}y^2(3y - a^2)(a^2 - y^2)^{\frac{1}{2}}dy; \\ \therefore A &= \frac{4}{a^4} \int_0^a (3y^2 - a^2)(a^2 - y^2)^{\frac{1}{2}}y^2dy = \frac{1}{8}a^2\pi. \end{aligned}$$

The equation of the curve JEK , is $x^{\frac{2}{3}} + y^{\frac{2}{3}} = (2a)^{\frac{2}{3}}$. Its length is $3a$, $JE = a$, and $KE = 2a$. Being the envelope of BL it is easily described, and hence furnishes a convenient means of describing the required curve.

[Mr. Seitz has also given a very elegant solution of this problem. He employs polar coordinates and obtains for the equation of the curve

$$\rho = \frac{8a \sin \theta \{ \sqrt{[(1 - \sin \theta)(1 + 3 \sin \theta)]} \pm \sqrt{[(1 + \sin \theta)(1 - 3 \sin \theta)]} \}}{\{ \sqrt{[(1 + \sin \theta)(1 + 3 \sin \theta)]} \pm \sqrt{[(1 - \sin \theta)(1 - 3 \sin \theta)]} \}^2}$$

and for length and area, respectively,

$$\begin{aligned} s &= 4 \int_0^{\frac{2}{3}a\sqrt{3}} \frac{\rho d\rho}{(\rho^2 - p^2)^{\frac{1}{2}}} + 4 \int_a^{\frac{2}{3}a\sqrt{3}} \frac{\rho d\rho}{(\rho^2 - p^2)^{\frac{1}{2}}} = a(4\sqrt{2} + \sin^{-1}\frac{1}{3}), \\ A &= 2 \int_a^{\frac{2}{3}a\sqrt{3}} \frac{p\rho d\rho}{(\rho^2 - p^2)^{\frac{1}{2}}} - 2 \int_0^{\frac{2}{3}a\sqrt{3}} \frac{p\rho d\rho}{(\rho^2 - p^2)^{\frac{1}{2}}} = \frac{1}{8}\pi a^2. \end{aligned}$$